

FOR KIDS AND TEENS WITH ENTHESITIS-RELATED  
ARTHRITIS OR PSORIATIC ARTHRITIS



**HELP YOUR CHILD  
FIND REAL RELIEF**

Patient portrayal.  
Individual results may vary.

In a clinical trial of kids and teens with enthesitis-related arthritis (ERA) or juvenile psoriatic arthritis (JPsA) taking COSENTYX® (secukinumab), those with ERA had a 53% reduced risk of flares, and those with JPsA had an 85% reduced risk of flares.

#### **INDICATION**

COSENTYX is a prescription medicine used to treat people 4 years of age and older with active enthesitis-related arthritis and people 2 years of age and older with active psoriatic arthritis.

#### **IMPORTANT SAFETY INFORMATION**

Do not use COSENTYX if you have had a severe allergic reaction to secukinumab or any of the other ingredients in COSENTYX. See the Medication Guide for a complete list of ingredients.

Please see additional Important Safety Information throughout brochure. Tap here for full [Prescribing Information](#), including [Medication Guide](#).

Visit [COSENTYX.com/juvenile-idiopathic-arthritis](https://www.cosentyx.com/juvenile-idiopathic-arthritis) for more information.



## KIDS AND TEENS WITH ERA AND JP<sub>s</sub>A HAVE A DIFFERENT OPTION

Juvenile idiopathic arthritis (JIA) is the most common form of Juvenile Arthritis (JA) in kids and teens. An overactive immune system is believed to ultimately cause the painful, swollen joints that come with JIA. There are 7 different types of JIA, and each type has its own symptoms. COSENTYX® (secukinumab) is the only medicine FDA approved to treat 2 types of JIA: ERA and JP<sub>s</sub>A.

**Read more to see why COSENTYX is the most prescribed medicine of its kind.\***

\*Across all indications combined.

### IMPORTANT SAFETY INFORMATION (continued) What is the most important information I should know about COSENTYX?

COSENTYX is a medicine that affects your immune system. COSENTYX may increase your risk of having serious side effects such as:

#### Infections

COSENTYX may lower the ability of your immune system to fight infections and may increase your risk of infections. Some people have died from these infections.

- Your doctor should check you for tuberculosis (TB) before starting treatment with COSENTYX.
- If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with COSENTYX and during treatment with COSENTYX.
- Your doctor should watch you closely for signs and symptoms of TB during treatment with COSENTYX.  
**Do not use COSENTYX if you have an active TB infection.**

Please see additional Important Safety Information throughout brochure. Tap here for full [Prescribing Information](#), including [Medication Guide](#).

Visit [COSENTYX.com/juvenile-idiopathic-arthritis](https://www.cosentyx.com/juvenile-idiopathic-arthritis) for more information. 3



## UNDERSTAND ERA & JPSA, THE TWO MOST COMMON FORMS OF JA

### WHAT IS ERA?

Enthesitis-related arthritis (ERA) is a type of juvenile idiopathic arthritis (JIA) that **affects the tissue where the muscles, ligaments, or tendons meet the bone** (entheses). Symptoms may include swelling, joint pain, and stiffness at the hips, knees, and feet. The fingers, elbows, pelvis, chest, and lower back can also be affected.

### WHAT IS JPSA?

Juvenile psoriatic arthritis (JPsA) is a type of JIA that **may include symptoms of both arthritis and plaque psoriasis**. Arthritis symptoms can show up before skin symptoms and may affect 1 or more joints, often in the wrists, ankles, fingers, or toes. Psoriasis can appear as a scaly rash behind the ears, on the eyelids, elbows, knees, belly button, or scalp.

## RECOGNIZE THE SIGNS OF A FLARE IN ERA & JPSA

When talking to a doctor about your child's symptoms, your doctor may refer to them as a flare. A flare is when your child's symptoms, involving at least 2 joints, become worse without much improvement. A flare can look different depending on the type of JIA and can last days or even months. It's important to minimize flares because they can have lasting effects.



### SYMPTOMS OF A FLARE MAY INCLUDE:

- Fatigue
- Increased stiffness that lasts for more than a few minutes
- Increased pain and swelling in joints
- Impacted range of motion
- Difficulty with usual activities
- Flu-like symptoms


### IMPORTANT SAFETY INFORMATION (continued)

#### Before starting COSENTYX®, tell your doctor if you:

- are being treated for an infection
- have an infection that does not go away or that keeps coming back
- have TB or have been in close contact with someone with TB
- think you have an infection or have symptoms of an infection such as: fevers, sweats, or chills; muscle aches; cough; shortness of breath; blood in your phlegm; weight loss; warm, red, or painful skin or sores on your body; diarrhea or stomach pain; burning when you urinate or urinate more often than normal

Please see additional Important Safety Information throughout brochure. Tap here for full [Prescribing Information](#), including [Medication Guide](#).

Visit [COSENTYX.com/juvenile-idiopathic-arthritis](https://www.cosentyx.com/juvenile-idiopathic-arthritis) for more information. 5



## COSENTYX HAS PROVEN EXPERIENCE

When you're considering COSENTYX<sup>®</sup> (secukinumab) as a treatment for your child's enthesitis-related arthritis (ERA) or juvenile psoriatic arthritis (JPsA), you should know that COSENTYX is the most prescribed medicine of its kind,<sup>\*</sup> and is approved to treat pediatric plaque psoriasis.

- **Over 3.7 million COSENTYX prescriptions** have been filled in the United States<sup>\*</sup>
- **Consistent safety profile** since its launch in 2015
- **No blood tests or routine lab monitoring** required during treatment with COSENTYX
- **Studied for more than 16 years** in dozens of clinical trials.<sup>\*</sup> More trials are ongoing as we stay committed to studying COSENTYX.

<sup>\*</sup>Across all indications combined.

### IMPORTANT SAFETY INFORMATION (continued)

**After starting COSENTYX, call your doctor right away if you have any signs of infection listed on page 5.** Do not use COSENTYX if you have any signs of infection unless you are instructed to by your doctor.

### What are the possible side effects of COSENTYX?

**COSENTYX may cause serious side effects, including:**

#### Serious allergic reactions

Serious allergic reactions can occur. Get emergency medical help right away if you get any of the following symptoms: feeling faint; swelling of your face, eyelids, lips, mouth, tongue, or throat; trouble breathing or throat tightness; chest tightness; skin rash or hives (red, itchy bumps).

**If you have a severe allergic reaction, do not give another injection of COSENTYX.**




Patient portrayal.  
Not for use in dogs.




## REAL RELIEF FROM 2 TYPES OF JUVENILE ARTHRITIS (JA)

In a clinical trial of kids and teens with enthesitis-related arthritis (ERA) or juvenile psoriatic arthritis (JPsA) taking COSENTYX, when results were reported at 2 years:

Those with ERA had



Those with JPsA had



A flare is when your child's symptoms become worse without much improvement, and involves at least 2 joints.

## SYMPTOM IMPROVEMENT IN AS EARLY AS 12 WEEKS

In a clinical trial of kids and teens taking COSENTYX<sup>®</sup> (secukinumab), 6 key areas were observed. Most (91% with JPsA and 85% with ERA) showed at least a 30% improvement in 3 or more of the 6 key areas, with worsening in no more than 1 key area, after 12 weeks of treatment.

- The doctor's assessment of disease activity
- The parent's or child's assessment of overall well-being
- Difficulty doing daily activities
- Number of joints affected by active arthritis
- Number of joints that are hard to move
- Lab test that measures inflammation

## HOW DOES COSENTYX WORK?

Having ERA or JPsA means your child has an overactive immune system. COSENTYX is a biologic treatment that works within the body's immune system to target and block a molecule called IL-17A, which may play a role in causing inflammation throughout the body when it is overproduced. Blocking IL-17A may help reduce the inflammation causing the joint pain and stiffness your child feels.

## WHAT IS A BIOLOGIC?

A biologic is a protein-based medication that is made from living sources and is taken by injection or infusion.

**TALK TO YOUR CHILD'S RHEUMATOLOGIST ABOUT COSENTYX TODAY.**

## IMPORTANT SAFETY INFORMATION (continued)

### Inflammatory bowel disease

New cases of inflammatory bowel disease or "flare-ups" can happen with COSENTYX, and can sometimes be serious. If you have inflammatory bowel disease (ulcerative colitis or Crohn's disease), tell your doctor if you have worsening disease symptoms during treatment with COSENTYX or develop new symptoms of stomach pain or diarrhea.

Please see additional Important Safety Information throughout brochure. Tap here for full [Prescribing Information](#), including [Medication Guide](#).

Visit [COSENTYX.com/juvenile-idiopathic-arthritis](https://www.cosentyx.com/juvenile-idiopathic-arthritis) for more information. 9

## GETTING YOUR CHILD STARTED WITH COSENTYX

Your child's doctor will prescribe the dose of COSENTYX® (secukinumab) that's right for your child based on their weight. Your child will take COSENTYX once a week for 5 weeks, then just once a month afterward.

- If your child weighs 33 lbs or more but less than 110 lbs, the recommended dose is 75 mg.
- If your child weighs 110 lbs or more, the recommended dose is 150 mg.

### COSENTYX REQUIRES JUST 1 INJECTION FOR EACH DOSE:

- A prefilled syringe for a 75 mg dose.
- A prefilled syringe or Sensoready® Pen for a 150 mg dose.

### IMPORTANT SAFETY INFORMATION (continued)

**Severe skin reactions that look like eczema** can happen during treatment with COSENTYX from days to months after your first dose and can sometimes lead to hospitalization. Your doctor may temporarily stop treatment with COSENTYX if you develop severe skin reactions. Tell your doctor if you have any of the following signs or symptoms: redness or rash; itching; small bumps or patches; your skin is dry or feels like leather; blisters on the hands or feet that ooze or become crusty or skin peeling.


## CONVENIENT ONCE-A-MONTH DOSING

### TO START

Your doctor will prescribe your child 5 weekly doses.\*

### TO CONTINUE

After that, **your child will only need to take COSENTYX once a month**, which means fewer interruptions in your daily routine.



\*Monthly dose equals 1 dose every 4 weeks.

### IMPORTANT SAFETY INFORMATION (continued)

**The most common side effects of COSENTYX include:** cold symptoms, diarrhea, and upper respiratory tract infections.

These are not all of the possible side effects of COSENTYX. Call your doctor for medical advice about side effects.

Please see additional Important Safety Information throughout brochure. Tap here for full [Prescribing Information](#), including [Medication Guide](#).

Visit [COSENTYX.com/juvenile-idiopathic-arthritis](https://www.cosentyx.com/juvenile-idiopathic-arthritis) for more information. 11



## GET COMFORTABLE TAKING COSENTYX

Your child's healthcare provider should show you how to prepare and inject COSENTYX<sup>®</sup> (secukinumab) before you use it for the first time. Please remember that your child should not inject themselves. Talk to your child's doctor if you have any questions.

For injection resources, visit [COSENTYX.com//kids-and-teens/how-to-inject-cosentyx](https://www.cosentyx.com/kids-and-teens/how-to-inject-cosentyx).

**TAP HERE TO VIEW SUPPLEMENTAL ▶  
INJECTION DEMONSTRATION VIDEOS.**



Patient portrayal.  
Individual results may vary.

## CREATE A ROUTINE

Real people taking COSENTYX have created personal routines around taking their injection. Routines can help ease the process and provide comfort as your child gets used to injections. Here are some ways to create an injection-time routine for your child:

- Find a peaceful spot in your home
- Listen to your child's favorite song
- Celebrate with a post-shot treat

### IMPORTANT SAFETY INFORMATION (continued)

**Before using COSENTYX, tell your doctor if you:**

- have any of the conditions or symptoms listed on page 5 for infections.
- have inflammatory bowel disease (Crohn's disease or ulcerative colitis).
- are allergic to latex. The needle cap on the COSENTYX Sensoready<sup>®</sup> pen, and 150 mg/mL and 75 mg/0.5 mL prefilled syringes contains latex.
- have recently received or are scheduled to receive an immunization (vaccine). People who take COSENTYX **should not** receive live vaccines. Children should be brought up to date with all vaccines before starting COSENTYX.
- have any other medical conditions and all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of your medicines to show your healthcare provider and pharmacist when you get a new medicine.
- are pregnant or plan to become pregnant. It is not known if COSENTYX can harm your unborn baby. You and your doctor should decide if you will use COSENTYX.
- are breastfeeding or plan to breastfeed. It is not known if COSENTYX passes into your breast milk.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout brochure. Tap here for full [Prescribing Information](#), including [Medication Guide](#).

Visit [COSENTYX.com/juvenile-idiopathic-arthritis](https://www.cosentyx.com/juvenile-idiopathic-arthritis) for more information.

## COSENTYX® CONNECT IS HERE FOR YOU

**COSENTYX® Connect is here throughout your treatment.** When you sign up, you'll have access to a full range of services and support to make your experience with COSENTYX® (secukinumab) as easy, affordable, and convenient as possible.



**Dedicated COSENTYX® Connect Team to help you get started**



### Co-pay\* offer

If your child has private insurance and is otherwise eligible. Subject to annual benefit limit. Additional limitations apply.

COVERED  
*until you're*  
COVERED

**Up to 2 years of COSENTYX for free** with the Covered Until You're Covered program† if your child's insurance coverage is denied



**Help with navigating insurance coverage**



**Personalized resources** via mail, e-mail, and text updates



**Injection training resources**

**TAP HERE TO SIGN UP. ►**



Patient portrayal.

\*Limitations apply. Valid only for those with private insurance. Program provides up to \$16,000 annually for the cost of COSENTYX and up to \$150 per infusion (up to \$1,950 annually) for the cost of administration. Co-pay support for infusion administration cost not available in Rhode Island or Massachusetts. Offer not valid under Medicare, Medicaid, or any other federal or state program. Novartis reserves the right to rescind, revoke, or amend this program without notice. See complete Terms & Conditions for details.

†The **Covered Until You're Covered Program** is available for COSENTYX (secukinumab) subcutaneous injection only. Eligible patients must have commercial insurance, a valid prescription for COSENTYX, and a denial of insurance coverage based on a prior authorization request. Program requires the submission of an appeal of the coverage denial within the first 90 days of enrollment in order to remain eligible. Program provides COSENTYX for free to eligible patients for up to two years, or until they receive insurance coverage approval, whichever occurs earlier. A valid prescription consistent with FDA-approved labeling is required. Program is not available to patients whose medications are reimbursed in whole or in part by Medicare, Medicaid, TRICARE, or any other federal or state program. Patients may be asked to reverify insurance coverage status during the course of the program. No purchase necessary. Program is not health insurance, nor is participation a guarantee of insurance coverage. Limitations may apply. Enrolled patients awaiting coverage for COSENTYX after two years may be eligible for a limited Program extension. Novartis Pharmaceuticals Corporation reserves the right to rescind, revoke, or amend this Program without notice.

Please see additional Important Safety Information throughout brochure. Tap here for full [Prescribing Information](#), including [Medication Guide](#).

Visit [COSENTYX.com/juvenile-idiopathic-arthritis](https://www.cosentyx.com/juvenile-idiopathic-arthritis) for more information. 15



## 3 EASY STEPS TO GET STARTED WITH COSENTYX

Choosing a new treatment is a big step, and once you and your child's doctor have decided that COSENTYX<sup>®</sup> (secukinumab) is right for your child, here is some information for you to consider to get going.



### 1 Sign Up Today

To enroll in the COSENTYX<sup>®</sup> Connect program and start receiving free resources, call **1-844-COSENTYX (1-844-267-3689)** or visit [COSENTYX.com](https://www.cosentyx.com).



### 2 Get COSENTYX Covered

Your COSENTYX<sup>®</sup> Connect Team Member will work with you and your healthcare provider to help get your child's COSENTYX prescription approved. You may be eligible for some of the saving options mentioned on page 14.



### 3 Get COSENTYX Delivered

COSENTYX is sent to you from a specialty pharmacy because it's a biologic medication, which means it requires special handling and must be refrigerated. **There are specific steps you'll have to take to arrange delivery.** When you enroll in COSENTYX<sup>®</sup> Connect you'll have a dedicated **COSENTYX<sup>®</sup> Connect Team Member to help guide you through every step of the process.** Make sure you save the specialty pharmacy phone number since they'll need to call you to schedule delivery of your COSENTYX right to your door.



Patient portrayal.

Please see additional Important Safety Information throughout brochure. Tap here for full [Prescribing Information](#), including [Medication Guide](#).

Visit [COSENTYX.com/juvenile-idiopathic-arthritis](https://www.cosentyx.com/juvenile-idiopathic-arthritis) for more information. 17



**QUESTIONS TO  
ASK YOUR CHILD'S DOCTOR?  
WRITE THEM HERE**

Lined area for writing questions to ask a child's doctor.

Please see additional Important Safety Information throughout brochure.  
Tap here for full Prescribing Information, including Medication Guide.  
Visit [COSENTYX.com/juvenile-idiopathic-arthritis](http://COSENTYX.com/juvenile-idiopathic-arthritis) for more information. 19





**Cosentyx**<sup>®</sup>  
(secukinumab)



Patient portrayal.  
Individual results may vary.

## ASK YOUR DOCTOR ABOUT COSENTYX

- COSENTYX<sup>®</sup> (secukinumab) is the **only medicine FDA approved to treat enthesitis-related arthritis (ERA) and juvenile psoriatic arthritis (JPsA)**
- Well-studied safety profile with **once-monthly dosing**\*

\*After starting with 5 weekly doses.

**TAP HERE TO SIGN UP FOR MORE INFORMATION ABOUT COSENTYX. ►**



Please see additional Important Safety Information throughout brochure. Tap here for full [Prescribing Information](#), including [Medication Guide](#).

Visit [COSENTYX.com/juvenile-idiopathic-arthritis](https://www.cosentyx.com/juvenile-idiopathic-arthritis) for more information.



**Novartis Pharmaceuticals Corporation**  
East Hanover, New Jersey 07936-1080  
© 2024 Novartis