


# HERE'S THE PLAN FOR YOUR NEXT DOCTOR APPOINTMENT

Having an informed conversation with your doctor and asking the right questions can make all the difference.

## 1 Mark the condition you're most concerned about

- Psoriatic arthritis (PsA)
- Plaque psoriasis (PsO)
- Ankylosing spondylitis (AS)
- Non-radiographic axial spondyloarthritis (nr-axSpA)
- Hidradenitis Suppurativa (HS)



## 2 Circle your most affected areas

Use the body outline to highlight your most affected areas.

## 4 Talk to your doctor about how your condition is affecting you

Write down any and all ways that your condition affects your day-to-day life.

---

---

## 5 Circle the face that best expresses your level of satisfaction with your current treatment



Very Unsatisfied



Unsatisfied



Indifferent



Satisfied



Completely Satisfied

- I'm not currently treating my condition.

**Be sure to discuss your treatment goals with your doctor during every visit.**

### Need help getting the conversation started about COSENTYX®? Ask these questions:

1. What makes COSENTYX a different type of biologic?

2. Do you think COSENTYX is right for me?

3. What are the benefits and risks of treating with COSENTYX?

**Take notes on your conversation to help you remember what you learned. For additional space, turn to page 3.**

---

---

### INDICATIONS

COSENTYX® (secukinumab) is a prescription medicine used to treat:

- people 6 years of age and older with moderate to severe plaque psoriasis (PsO) that involves large areas or many areas of the body, and who may benefit from taking injections or pills (systemic therapy) or phototherapy (treatment using ultraviolet or UV light alone or with systemic therapy)
- people 2 years of age and older with active psoriatic arthritis (PsA)
- adults with active ankylosing spondylitis (AS)
- adults with active non-radiographic axial spondyloarthritis (nr-axSpA) and objective signs of inflammation
- people 4 years of age and older with active enthesitis-related arthritis (ERA)
- adults with moderate to severe hidradenitis suppurativa (HS)

### IMPORTANT SAFETY INFORMATION

Do not use COSENTYX if you have had a severe allergic reaction to secukinumab or any of the other ingredients in COSENTYX. See the Medication Guide for a complete list of ingredients.

**Please see full Prescribing Information, including Medication Guide.**

Visit [COSENTYX.com](http://COSENTYX.com) for more information.



## IMPORTANT SAFETY INFORMATION (cont)

### What is the most important information I should know about COSENTYX?

COSENTYX® (secukinumab) is a medicine that affects your immune system. COSENTYX may increase your risk of having serious side effects such as:

#### Infections

COSENTYX may lower the ability of your immune system to fight infections and may increase your risk of infections. Some people have died from these infections.

- Your doctor should check you for tuberculosis (TB) before starting treatment with COSENTYX.
- If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with COSENTYX and during treatment with COSENTYX.
- Your doctor should watch you closely for signs and symptoms of TB during treatment with COSENTYX. **Do not use COSENTYX if you have an active TB infection.**

#### Before starting COSENTYX, tell your doctor if you:

- are being treated for an infection
- have an infection that does not go away or that keeps coming back
- have TB or have been in close contact with someone with TB
- think you have an infection or have symptoms of an infection such as: fevers, sweats, or chills; muscle aches; cough; shortness of breath; blood in your phlegm; weight loss; warm, red, or painful skin or sores on your body; diarrhea or stomach pain; burning when you urinate or urinate more often than normal

**After starting COSENTYX, call your doctor right away if you have any signs of infection listed above.** Do not use COSENTYX if you have any signs of infection unless you are instructed to by your doctor.

### What are the possible side effects of COSENTYX?

#### COSENTYX may cause serious side effects, including:

##### Serious allergic reactions

Serious allergic reactions can occur. Get emergency medical help right away if you get any of the following symptoms: feeling faint; swelling of your face, eyelids, lips, mouth, tongue, or throat; trouble breathing or throat tightness; chest tightness; skin rash or hives (red, itchy bumps).

**If you have a severe allergic reaction, do not give another injection of COSENTYX.**

##### Inflammatory bowel disease

New cases of inflammatory bowel disease or "flare-ups" can happen with COSENTYX, and can sometimes be serious. If you have inflammatory bowel disease (ulcerative colitis or Crohn's disease), tell your doctor if you have worsening disease symptoms during treatment with COSENTYX or develop new symptoms of stomach pain or diarrhea.

**Severe skin reactions that look like eczema** can happen during treatment with COSENTYX from days to months after your first dose and can sometimes lead to hospitalization. Your doctor may temporarily stop treatment with COSENTYX if you develop severe skin reactions. Tell your doctor if you have any of the following signs or symptoms: redness or rash; itching; small bumps or patches; your skin is dry or feels like leather; blisters on the hands or feet that ooze or become crusty or skin peeling.

**The most common side effects of COSENTYX include:** cold symptoms, diarrhea, and upper respiratory tract infections.

These are not all of the possible side effects of COSENTYX. Call your doctor for medical advice about side effects.

#### Before using COSENTYX, tell your doctor if you:

- have any of the conditions or symptoms listed in the previous column for infections.
- have inflammatory bowel disease (Crohn's disease or ulcerative colitis)
- are allergic to latex. The needle cap on the COSENTYX Sensoready® pen, and 150 mg/mL and 75 mg/0.5 mL prefilled syringes contains latex.
- have recently received or are scheduled to receive an immunization (vaccine). People who take COSENTYX **should not** receive live vaccines. Children should be brought up to date with all vaccines before starting COSENTYX.
- have any other medical conditions and all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of your medicines to show your healthcare provider and pharmacist when you get a new medicine.
- are pregnant or plan to become pregnant. It is not known if COSENTYX can harm your unborn baby. You and your doctor should decide if you will use COSENTYX.
- are breastfeeding or plan to breastfeed. It is not known if COSENTYX passes into your breast milk.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

**Please see full [Prescribing Information](#), including [Medication Guide](#).**

Visit [COSENTYX.com](http://COSENTYX.com) for more information.



