



## Honey Mustard Glazed Salmon with Roasted Vegetables

By Chef Michelle Bernstein



*"It's the perfect time to roast some beautiful vegetables and eat them with your favorite fish, including salmon, which we all know is easy to find and rich with all kinds of wonderful nutrients."*

Chef Michelle is an actual patient and has been compensated for her time. Individual results may vary.

### Ingredients:

- 4 (6–7 ounce) pieces salmon, skin on, center cut
- 3 heaping tablespoons grainy mustard
- 1 heaping tablespoon Dijon mustard
- 2 teaspoons balsamic vinegar
- 3 tablespoons honey
- 1 teaspoon garlic cloves, minced
- ½ teaspoon salt
- 2 bunches broccolini, cut into 6–8 long pieces each
- ½ acorn squash, seeds removed, sliced into ¼-inch thick wedges (with peel)
- ½ small red onion, cut into thick strips
- 2 cups Brussels sprout leaves
- Salt and pepper, to taste
- Olive oil for drizzling
- 1 fresh lemon
- Coarse sea salt, if desired
- Nonstick cooking spray

### Directions:



#### Step 1

Preheat your oven to 400 °F. Line a rimmed baking sheet with parchment paper or foil. Place acorn squash and red onions on the tray, drizzle with olive oil, and season with salt and pepper to taste. Bake for 8 minutes.



#### Step 2

Remove pan and add the broccolini and Brussels sprout leaves, drizzle with a little olive oil, and season with salt and pepper. Cook another 8–10 minutes.



#### Step 3

When the vegetables are cooked, top them with some zest from the lemon. While the vegetables roast, line a second rimmed baking sheet with parchment paper and spray with nonstick cooking spray. Set aside.



#### Step 4

Whisk together the 2 mustards, balsamic vinegar, honey, minced garlic, and salt. Place the salmon, skin side down, on the baking sheet. Generously spread the glaze over the salmon filets.



#### Step 5

Bake the salmon at 400°F for 6–8 minutes or to your desired doneness. Top the salmon with a little coarse sea salt, if desired. Serve the salmon filets, discarding the skin, with the roasted vegetables.

## TIPS TO HELP WITH YOUR NEXT DOCTOR'S APPOINTMENT

Here are 4 tips that can help you reflect on your psoriatic arthritis symptoms and discuss treatment options with your doctor.

### TIP 1: Reflect on Your Symptoms

Take a moment to think about how psoriatic arthritis is impacting your day-to-day life. Where are you experiencing symptoms like joint pain, swelling, and tenderness, and how painful are they?

Which daily activities are affected by your psoriatic arthritis symptoms? Check all that apply.

Exercising ☐ Running Errands ☐ Getting out of Bed ☐ Other ☐  
Gripping ☐ Preparing Meals ☐ Getting Dressed ☐

### TIP 2: Reflect on Your Treatment

Circle the face that best expresses your level of satisfaction with your current treatment.



Very  
Unsatisfied



Unsatisfied



Indifferent



Satisfied



Completely  
Satisfied

☐ I'm not currently treating my psoriatic arthritis.

### TIP 3: Speak Up

Now that you've spent time reflecting, make sure you tell your doctor about your symptoms and how they're really making you feel.

### TIP 4: Advocate for Yourself

In addition to speaking up, explore your treatment options with your doctor. Ask the questions below to find out if COSENTYX® is right for you:

1. What makes COSENTYX different?
2. Do you think COSENTYX is right for me?
3. What are the benefits and risks of treating with COSENTYX?

#### INDICATIONS

COSENTYX® (secukinumab) is a prescription medicine used to treat:

- people 6 years of age and older with moderate to severe plaque psoriasis (PsO) that involves large areas or many areas of the body, and who may benefit from taking injections or pills (systemic therapy) or phototherapy (treatment using ultraviolet or UV light alone or with systemic therapy)
- people 2 years of age and older with active psoriatic arthritis (PsA)
- adults with active ankylosing spondylitis (AS)
- adults with active non-radiographic axial spondyloarthritis (nr-axSpA) and objective signs of inflammation
- people 4 years of age and older with active enthesitis-related arthritis (ERA)
- adults with moderate to severe hidradenitis suppurativa (HS)

#### IMPORTANT SAFETY INFORMATION

Do not use COSENTYX if you have had a severe allergic reaction to secukinumab or any of the other ingredients in COSENTYX. See the Medication Guide for a complete list of ingredients.

**Please see additional Important Safety Information on the next page.**

**Click here for full Prescribing Information, including Medication Guide.**

Visit [COSENTYX.com](https://www.cosentyx.com) for more information.

## IMPORTANT SAFETY INFORMATION (cont)

### What is the most important information I should know about COSENTYX?

COSENTYX is a medicine that affects your immune system. COSENTYX may increase your risk of having serious side effects such as:

#### Infections

COSENTYX may lower the ability of your immune system to fight infections and may increase your risk of infections. Some people have had serious infections while taking COSENTYX, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses. Some people have died from these infections.

- Your doctor should check you for TB before starting treatment with COSENTYX.
- If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with COSENTYX and during treatment with COSENTYX.
- Your doctor should watch you closely for signs and symptoms of TB during treatment with COSENTYX. **Do not take COSENTYX if you have an active TB infection.**

#### Before starting COSENTYX, tell your doctor if you:

- are being treated for an infection
- have an infection that does not go away or that keeps coming back
- have TB or have been in close contact with someone with TB
- think you have an infection or have symptoms of an infection such as: fevers, sweats, or chills; muscle aches; cough; shortness of breath; blood in your phlegm; weight loss; warm, red, or painful skin or sores on your body; diarrhea or stomach pain; burning when you urinate or urinate more often than normal

**After starting COSENTYX, call your doctor right away if you have any signs of infection listed above.** Do not use COSENTYX if you have any signs of infection unless you are instructed to by your doctor.

#### What are the possible side effects of COSENTYX?

COSENTYX may cause serious side effects, including:

#### Serious allergic reactions

Serious allergic reactions can occur. Get emergency medical help right away if you get any of the following symptoms: feeling faint; swelling of your face, eyelids, lips, mouth, tongue, or throat; trouble breathing or throat tightness; chest tightness; skin rash or hives (red, itchy bumps).

**If you have a severe allergic reaction, do not give another injection of COSENTYX.**

#### Inflammatory bowel disease

New cases of inflammatory bowel disease or “flare-ups” can happen with COSENTYX, and can sometimes be serious. If you have inflammatory bowel disease (ulcerative colitis or Crohn's disease), tell your doctor if you have worsening disease symptoms during treatment with COSENTYX or develop new symptoms of stomach pain or diarrhea.

**Severe skin reactions that look like eczema** can happen during treatment with COSENTYX from days to months after your first dose and can sometimes lead to hospitalization. Your doctor may temporarily stop treatment with COSENTYX if you develop severe skin reactions. Tell your doctor if you have any of the following signs or symptoms: redness or rash; itching; small bumps or patches; your skin is dry or feels like leather; blisters on the hands or feet that ooze or become crusty or skin peeling.

**The most common side effects of COSENTYX include:** cold symptoms, diarrhea, and upper respiratory tract infections.

These are not all of the possible side effects of COSENTYX. Call your doctor for medical advice about side effects.

#### Before using COSENTYX, tell your doctor if you:

- have any of the conditions or symptoms listed above for infections.
- have inflammatory bowel disease (Crohn's disease or ulcerative colitis).
- are allergic to latex. The needle cap on the COSENTYX Sensoready<sup>®</sup> pen, and 150 mg/mL and 75 mg/0.5 mL prefilled syringes contains latex.
- have recently received or are scheduled to receive an immunization (vaccine). People who take COSENTYX **should not** receive live vaccines. Children should be brought up to date with all vaccines before starting COSENTYX.
- have any other medical conditions and all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of your medicines to show your healthcare provider and pharmacist when you get a new medicine.
- are pregnant or plan to become pregnant. It is not known if COSENTYX can harm your unborn baby. You and your doctor should decide if you will use COSENTYX.
- are breastfeeding or plan to breastfeed. It is not known if COSENTYX passes into your breast milk.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

Please see full **Prescribing Information**, including **Medication Guide**.

Visit [COSENTYX.com](http://COSENTYX.com) for more information.



Novartis Pharmaceuticals Corporation  
East Hanover, New Jersey 07936-1080

© 2024 Novartis

8/24

FA-11241454

