USE THIS CHECKLIST WHEN GETTING STARTED ON COSENTYX® (secukinumab)



SAVE YOUR SPECIALTY PHARMACY CONTACT IN YOUR PHONE

- Remember to ask your doctor for your specialty pharmacy name and phone number
- Your healthcare provider, insurance carrier, specialty pharmacy, and Novartis Patient Support will all work together to help get your prescription processed, approved, and delivered



PICK UP THE PHONE WHEN THE PHARMACY CALLS

Your specialty pharmacy will call to schedule your delivery



If the pharmacy calls from a 1-800 number you don't recognize, make sure you answer or call the pharmacy back right away so your delivery isn't delayed. You'll also want to save that number for future use.

 If you don't hear from the specialty pharmacy after a few weeks, please call the pharmacy or your doctor's office to follow up

| Specialty Pharmacy Name: | P _X |
|--|---|
| Specialty Pharmacy Phone: | |
| Name of Medication: COSENTYX® (| (secukinumab) |
| Fill this out with your doctor, so yo about your dose. | ou're clear |
| My dose is: | |
| ☐ 75 mg (pediatric ☐ 150 mg dosing) | 300 mg (taken as two 150 mg injections or one 300 mg injection) |
| To start, your doctor will prescribe or just 1 dose once a month, based After that, you only need to take CC | on what's right for you. |
| If you have HS and continue to have may prescribe COSENTYX every 2 is given as 1 injection of 300 mg or 1 | weeks. Each 300-mg dos |

[‡]Monthly dose equals 1 dose every 4 weeks.



SIGN UP FOR NOVARTIS PATIENT SUPPORT EVEN BEFORE TAKING YOUR FIRST DOSE*

Get access to the full range of benefits to help you every step of the way





SIGN UP FOR NOVARTIS PATIENT SUPPORT BY VISITING <u>COSENTYX.COM/REGISTER</u> OR CALLING 1-844-COSENTYX (1-844-267-3689).

Watch a step-by-step video to help with getting started at COSENTYX.com/gettingstarted.

*Novartis Patient Support enrollment is optional and not required for getting started on treatment.

†For terms, conditions, and limitations related to Co-Pay Plus offer and Covered Until You're Covered also known as Bridge Program, please see the back of this page.



HERE'S HOW THE PROCESS WORKS





COSENTYX® (secukinumab) PRESCRIPTION

Your doctor sends the prescription to the specialty pharmacy of your choice, or to an insurance-mandated pharmacy.



Novartis Patient Support ENROLLMENT*

Sign up for Novartis Patient Support to get financial support, if eligible, and resources to help you every step of the way. Sign up by visiting **COSENTYX.com/register** or calling 1-844-COSENTYX (1-844-267-3689).



Once enrolled, you will receive a welcome call from your dedicated **Novartis Patient Support Team** Member. They will be calling from 1-844-267-3689, so be sure to pick up the phone.





INSURANCE COVERAGE AND SAVINGS OPTIONS

Your doctor, insurance carrier, specialty pharmacy, and Novartis Patient Support will all work together to

- · A specialty pharmacy needs to be identified
- Your prescription is covered by insurance
- Prior authorization (PA) is required Your insurance provider may require your doctor to provide additional information (ie, medical history, lab work, etc) before it will cover the medication.



Co-Pay Plus offer

PRIOR AUTHORIZATION APPROVED

Once you are approved, your specialty pharmacy will be notified to process delivery. Don't forget to sign up for Novartis Patient Support to receive \$0 Co-Pay Plus offer† if you are eligible.



PRIOR AUTHORIZATION DENIED

If your insurance is initially denied, don't stop there. You may be eligible to receive up to 2 years of FREE COSENTYX through our Covered Until You're Covered program, while we work with your healthcare provider for coverage. Ask your doctor for more information.





DELIVERY

COSENTYX is a biologic and must be delivered by a specialty pharmacy. Your Novartis Patient Support Team Member or your specialty pharmacy will be calling to ask for additional information and to schedule delivery.



10/25

Be sure to pick up the phone. It may be from an unfamiliar 1-800 number. Remember to save the specialty pharmacy number in your phone for future reference.



CONTINUED SUPPORT

With Novartis Patient Support, you'll get access to educational tools and support, injection training videos, a dedicated Novartis Patient Support Team member to assist you every step of the way, and inspirational emails and texts to help keep you going. We'll even check in from time to time via email, text, or phone.

*Novartis Patient Support enrollment is optional and not required for getting started on treatment.

*Limitations apply. Subject to annual co-pay benefit limit. Offer not valid under Medicare, Medicaid, or any other federal or state programs. Novartis reserves the right to rescind, revoke, or amend this program without notice. Additional limitations may apply. See complete Terms & Conditions at [https://www.cosentyx.com/all/treatment-cost#terms-and-conditions] for details.

*Covered Until You're Covered also known as Bridge Program applies to COSENTYX Subcutaneous Injection only. Eligible patients must have private insurance and a valid prescription for COSENTYX, and a prior

authorization, predetermination, or medical exception that has been denied. Program requires the submission of an appeal of the coverage denial within the first 90 days of enrollment to remain eligible. Program provides COSENTYX for free to eligible patients for up to 2 years, or until they receive insurance coverage approval, whichever occurs earlier. A valid prescription consistent with FDA-approved labeling is required. Program is not available to patients whose medications are reimbursed in whole or in part by Medicare, Medicaid, TRICARE, or any other federal or state program. Patients may be asked to reverify insurance coverage status during the course of the program. No purchase necessary. Program is not health insurance, nor is participation a guarantee of insurance coverage. Additional Limitations may apply. Novartis Pharmaceuticals Corporation reserves the right to rescind, revoke, or amend this Program without notice.

Please see Important Safety Information on page 3. Please see full Prescribing Information, including Medication Guide.



INDICATIONS

COSENTYX® (secukinumab) is a prescription medicine used to treat:

- people 6 years of age and older with moderate to severe plaque psoriasis (PsO) that involves large areas or many areas of the body, and who may benefit from taking injections or pills (systemic therapy) or phototherapy (treatment using ultraviolet or UV light alone or with systemic therapy)
- people 2 years of age and older with active psoriatic arthritis (PsA)
- adults with active ankylosing spondylitis (AS)
- adults with active non-radiographic axial spondyloarthritis (nr-axSpA) and objective signs of inflammation
- people 4 years of age and older with active enthesitis-related arthritis (ERA)
- adults with moderate to severe hidradenitis suppurativa (HS)

IMPORTANT SAFETY INFORMATION

Do not use COSENTYX if you have had a severe allergic reaction to secukinumab or any of the other ingredients in COSENTYX. See the Medication Guide for a complete list of ingredients.

What is the most important information I should know about COSENTYX?

COSENTYX is a medicine that affects your immune system. COSENTYX may increase your risk of having serious side effects such as:

Infections

COSENTYX may lower the ability of your immune system to fight infections and may increase your risk of infections. Some people have had serious infections while taking COSENTYX, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses. Some people have died from these infections.

- · Your doctor should check you for TB before starting treatment with COSENTYX.
- If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with COSENTYX and during treatment with COSENTYX.
- Your doctor should watch you closely for signs and symptoms of TB during treatment with COSENTYX. Do not use COSENTYX if you have an
 active TB infection.

Before starting COSENTYX, tell your doctor if you:

- · are being treated for an infection
- have an infection that does not go away or that keeps coming back
- · have TB or have been in close contact with someone with TB
- think you have an infection or have symptoms of an infection such as: fevers, sweats, or chills; muscle aches; cough; shortness of breath; blood
 in your phlegm; weight loss; warm, red, or painful skin or sores on your body; diarrhea or stomach pain; burning when you urinate or urinate more
 often than normal

After starting COSENTYX, call your doctor right away if you have any signs of infection listed above. Do not use COSENTYX if you have any signs of infection unless you are instructed to by your doctor.

What are the possible side effects of COSENTYX?

COSENTYX may cause serious side effects, including:

Serious allergic reactions

Serious allergic reactions can occur. Get emergency medical help right away if you get any of the following symptoms: feeling faint; swelling of your face, eyelids, lips, mouth, tongue, or throat; trouble breathing or throat tightness; chest tightness; skin rash or hives (red, itchy bumps).

If you have a severe allergic reaction, do not give another injection of COSENTYX.

Inflammatory bowel disease

New cases of inflammatory bowel disease or "flare-ups" can happen with COSENTYX, and can sometimes be serious. If you have inflammatory bowel disease (ulcerative colitis or Crohn's disease), tell your doctor if you have worsening disease symptoms during treatment with COSENTYX or develop new symptoms of stomach pain or diarrhea.

Severe skin reactions that look like eczema can happen during treatment with COSENTYX from days to months after your first dose and can sometimes lead to hospitalization. Your doctor may temporarily stop treatment with COSENTYX if you develop severe skin reactions. Tell your doctor if you have any of the following signs or symptoms: redness or rash; itching; small bumps or patches; your skin is dry or feels like leather; blisters on the hands or feet that ooze or become crusty or skin peeling.

The most common side effects of COSENTYX include: cold symptoms, diarrhea, and upper respiratory tract infections.

These are not all of the possible side effects of COSENTYX. Call your doctor for medical advice about side effects.

Before using COSENTYX, tell your doctor if you:

- · have any of the conditions or symptoms listed above for infections.
- have inflammatory bowel disease (Crohn's disease or ulcerative colitis).
- are allergic to latex. The needle cap on the COSENTYX Sensoready® pen, and 150 mg/mL and 75 mg/0.5 mL prefilled syringes contains latex.
- have recently received or are scheduled to receive an immunization (vaccine). People who take COSENTYX should not receive live vaccines.
 Children should be brought up to date with all vaccines before starting COSENTYX.
- have any other medical conditions and all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of your medicines to show your healthcare provider and pharmacist when you get a new medicine.
- are pregnant or plan to become pregnant. It is not known if COSENTYX can harm your unborn baby. You and your doctor should decide if you will use COSENTYX.
- · are breastfeeding or plan to breastfeed. It is not known if COSENTYX passes into your breast milk.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see full Prescribing Information, including Medication Guide.